Southampton Solent University

High- and Low-Load Resistance Training: Interpretation and Practical Application of Current Research Findings

Fisher, James; Steele, James; Smith, Dave

Published in:
Sports Medicine

DOI:
10.1007/s40279-016-0602-1

Published: 01/08/2016

Document Version
Peer reviewed version

Link to publication

Citation for pulished version (APA):
Fisher, J., Steele, J., & Smith, D. (2016). High- and Low-Load Resistance Training: Interpretation and Practical Application of Current Research Findings. *Sports Medicine*, *47*(3), 393-400. https://doi.org/10.1007/s40279-016-0602-1

General rights
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- Unless given explicit permission, you may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy
If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 10. Oct. 2026

High- and Low-load resistance training; Interpretation and Practical Application of Current Research findings

Current Opinion

James Fisher [1], James Steele [1], Dave Smith [2]

[1] Southampton Solent University, East Park Terrace, Southampton, UK

[2]Manchester Metropolitan University, Crewe, UK

Corresponding Author:

James Fisher

Southampton Solent University

East Park Terrace

Southampton

UK

Tel: +44 2382 013163

Email: James.Fisher@solent.ac.uk

Co-Author contact details:

James Steele (James.steele@solent.ac.uk)

Dave Smith (D.D.Smith@mmu.ac.uk)

Abstract word count: 190

Text only word count: 3321

Key Points

Current research is equivocal regarding the use of heavy- or light-loads for optimal strength and hypertrophic adaptations.

Misinterpretation of EMG amplitude, differing hypertrophic assessment methods (e.g. *in vivo* and *in vitro*) and unconsidered motor schema research might present reasons behind the differing adaptations reported.

Abstract

Our current state of knowledge regarding the load (lighter or heavier) lifted in resistance training programs to result in 'optimal' strength and hypertrophic adaptations is unclear. Despite this, position stands and recommendations are made based on, we propose, limited evidence to lift heavier weights. Here we discuss the state of evidence on the impact of load and how it, as a single variable, stimulates adaptations to take place and whether evidence for recommending heavier loads is available, well defined, currently correctly interpreted, or has been overlooked. Areas of discussion include electromyography amplitude, *in vivo* and *in vitro* methods of measuring hypertrophy, and motor schema and skill acquisition. The present piece clarifies to trainers and trainees the impact of these variables by discussing interpretation of synchronous and sequential motor unit recruitment and revisiting the size principle, poor agreement between whole muscle cross sectional area (CSA) and biopsy determined changes in myofibril CSA, and neural adaptations around task specificity. Our opinion is that the practical implications of being able to self-select external load might reduce the need for specific facility memberships, motivate older persons or those who might be less confident using heavy loads, and allow people to undertake home- or field-based resistance training intervention strategies which might ultimately improve exercise adherence.

50

51

52

1. Introduction

The role of load within resistance training is presently a hotly discussed topic in exercise science. Recent reviews have examined existing studies comparing the effects of different loads upon muscle function (e.g. strength and endurance) and hypertrophy. In these reviews some authors have suggested that, when resistance training is continued to momentary failure, essentially the same adaptations are possible with both heavy-(HL) or light-loads (LL) [1,2]. In contrast, others suggest that inclusion of specifically LL or HL may be necessary for optimising certain adaptations [3-6]. We propose that 'heavy' and 'light' loading systems exist on a spectrum and are individual based on subjectivity, however, for clarity, HL and LL have been operationally defined as >65% 1-repetition maximum (RM) and <60% 1RM, respectively [6]. A number of recent studies have been published examining both acute mechanistic differences resulting from difference in load in addition to studies comparing chronic changes in muscle function and hypertrophy. Unfortunately, we believe that some researchers may have inappropriately interpreted the data produced in these studies, with much of this surrounding incorrect inferences regarding motor unit (MU) recruitment in acute studies of electromyography (EMG), in addition to different methods of measuring both muscle function and hypertrophy. With this in mind, in the present piece we aim to discuss why different exercise scientists might have given contrasting recommendations by discussing the factors that should be considered in interpretation of research within this area.

2. Acute EMG Amplitude and the Size Principle

It is commonly accepted within the resistance training literature that recruitment of a MU is necessary in order for subsequent adaptation to occur [7]. Since discussions around optimal load for muscular adaptations are predicated upon the belief that complete

recruitment of MUs and thus muscle fibres is required for optimal adaptations, it is essential to consider acute EMG research within this area as well as briefly reconsider the size principle of motor unit recruitment. Recent studies have reported higher peak EMG amplitude for HL compared with LL [7,8] with one recent study showing increasing EMG amplitudes from 50% to 70% and to 90% 1RM [9]. From this the authors of these studies have inferred that LL do not maximally recruit all MU and as such HL are favourable for development of strength and hypertrophy. However, such recommendations may be founded upon an incorrect use and interpretation of EMG data relating to MU recruitment as well as a misapplication of the size principle.

For clarity, the size principle states that *"when the central nervous system recruits motor units for a specific activity it begins with the smallest, more easily excited, least powerful motor units and progresses to the larger, more difficult to excite, most powerful motor units to maintain or increase force"* [10,11]. However, as noted recently by Enoka and Duchateau [12], whilst EMG amplitude is influenced by MU recruitment strategies many continue to mistakenly infer MU recruitment from amplitude data. For example during a maximal voluntary contraction more MUs, including both those of a low or high threshold, will be activated and at increased frequencies in order to produce maximal force. As such the high MU recruitment would result in a higher EMG amplitude. In comparison, a sustained submaximal contraction would only recruit sufficient MUs to produce the necessary force; however, as those MUs fatigue other MUs would be recruited to replace them in sustaining the desired force. Indeed, during fatiguing contractions the threshold for recruitment of higher threshold MU is reduced permitting their subsequent recruitment [13] and MUs may 'cycle' (momentary de-recruitment and recruitment of different MU) during submaximal fatiguing contractions to reduce fatigue and maintain force [14]. Furthermore the 'muscle wisdom hypothesis' suggests that during sustained contractions motor unit discharge rate might

decrease due to optimizing the force output of motor units and protecting against peripheral conduction failure (Petrofsky & Phillips, 1985; Behm, 2004). Should this decrease in discharge rate occur, there would be a resultant decrease in signal amplitude (Garland & Gossen, 2002). As such, whilst HL would require more synchronous MU recruitment at greater frequencies (resulting in higher EMG amplitudes), sustained contractions to muscular failure with LL might ultimately recruit all MUs albeit sequentially (resulting in lower EMG amplitudes) rather than synchronously.

It should be noted that whether MU recruitment is ultimately similar between HL and LL remains a hypothesis to test empirically. Examination of this would require more advanced handling of EMG data such as spike-triggered averaging [15] or initial wavelet analysis followed by principal component classification of major frequency properties and optimization to tune wavelets to these frequencies [16]. Though acute mechanistic data cannot be used to infer chronic adaptations, studies such as these recent EMG amplitude comparisons of HL and LL are useful for generating hypotheses for examination in chronic training interventions. However, the hypotheses presented by the authors of these recent studies suggesting that HL may produce greater adaptations appear to stem from inappropriate interpretation of EMG amplitudes and consideration of the size principle.

3. Hypertrophic Adaptations

Common methods of measuring hypertrophy are *in vivo* (e.g. computed tomography [CT], magnetic resonance imaging [MRI] and ultrasound [UT]) and *in vitro* (e.g. muscle biopsy). Recent reviews have differed in their inclusion of studies using these methods with some opting to examine only *in vivo* measures of whole muscle hypertrophy [1] and others considering both *in vivo* and *in vitro* measures [5,6]. In fact, methods used to measure hypertrophy, the information they can provide and the strengths and weaknesses of both have

been discussed in light of these publications [17,18]. We acknowledge that whilst both *in vivo* and *in vitro* methods present useful information both offer very different information and that the two should be interpreted individually and carefully.

In both a recent review [5] and meta-analysis [6] of hypertrophy in response to HL and LL, resistance training studies utilising both muscle biopsy and *in vivo* methods were considered, and in the meta-analysis combined for analysis. The combination of *in vivo* and *in vitro* measures in this meta-analysis might have confounded the overall conclusions drawn in relation to other publications [1]. In support of this concern, a study by Mitchell et al. [19] included in the meta-analysis conducted both MRI and biopsy measures of hypertrophy in response to different resistance training loads and reported that relative increases appear greater for biopsy measures (mean = ~17-30% type I and ~16-18% type II; favouring low and high load conditions respectively in terms of effect size) compared to MRI (~7%; favouring the high load condition in terms of effect size). McCall et al. [20] have also reported differences between muscle biopsy and MRI methods in magnitude of CSA increase (mean =; biopsy = 10% type I fibre and 17% type II vs. 11.2% MRI). It is not clear from the meta-analysis method section how the authors dealt with the inclusion of the different outcome measures for hypertrophy used by Mitchell et al. [19], i.e. whether they were dealt with separately or combined. Indeed it has been noted [17] that in the earlier review [5] those studies using *in vivo* measures of whole muscle hypertrophy consistently showed no difference between HL and LL whereas the two *in vitro* studies using biopsies did show significantly greater gains for HL. Whilst ultimately still not statistically significant (p=0.076), it is unclear the degree to which the combination of methods influenced the results of this meta-analysis in favour of greater ESs for HL compared to LL ((mean ±SD) LL= 0.39 ±0.17; HL= 0.82 ±0.17). Figure 1 shows an adaptation of the forest plot originally published by Schoenfeld, et al. [6]. The dotted line shows the overall ES and clearly shows studies

151 which considered in vivo methods to the left of the dotted line, and studies which considered
152 in vitro methods to the right of the line, which we propose might have contaminated the
153 analyses and overall outcome.

154 The use of *in vitro* measures such as muscle biopsy permits the examination of many
155 important aspects of muscular adaptation including individual fibre typing, individual fibre
156 area, mitochondrial content, enzyme expression, capillarisation. Indeed it has been suggested
157 that fibre type specific adaptations may occur in response to HL or LL training [21] and,
158 though evidence is mixed at present as to whether this indeed occurs [19,22,23], biopsy
159 would be necessary to test this hypothesis further. Pertinent to hypertrophy as an outcome it
160 has been argued that a case could made for biopsy providing the most relevant information,
161 as individual fibre area can be determined, thus allowing differentiation between contractile
162 and non-contractile components [5]. However, it should be noted that evidence is equivocal
163 regarding the agreement between whole muscle CSA changes and biopsy-determined
164 changes in myofibril CSA, with some studies suggesting similar magnitude of relative change
165 [24,25] whereas others do not [26,27]. In fact, authors have actually agreed that *"it might be*
166 *true…that single fiber CSA data over-estimate whole muscle CSA"* [Burd, et al. 2013;
167 Schuenke, et al. 2013]. Methods exist to ensure sufficient tissue samples are obtained for
168 analysis using biopsy, yet, only a limited number of cells are assessed irrespective of method.
169 In this sense, variation in fibre characteristics and non-uniform growth along the length of a
170 muscle [28] provide notable limitations in attempting to extrapolate biopsy results to consider
171 whole muscle change [29]. However, measuring muscular adaptation using *in vivo* methods is
172 not without issues: different methods (MRI, CT, UT) can offer different information for both
173 individual and whole muscle groups including CSA, muscle thickness, muscle density,
174 architectural changes such as pennation angle, and changes in non-contractile components
175 such as intra-muscular adipose tissue. Again pertinent to the outcome of hypertrophy, even

consideration of whole muscle changes in CSA or muscle thickness may not be fully reflective of morphological adaptation. CSA may also include non-contractile components and so increases may not fully reflect muscular adaptations. Further and conversely, prior studies have reported lack of change in CSA yet significant increases in muscular density [30] in addition to disproportionate strength and CSA gains possibly being influenced by changes in muscle density [31].

In our opinion, the confounding factors discussed limit the integrity of any outcome data where analyses have combined these methods of measurement of hypertrophy. Furthermore, from a practical perspective, different outcomes may hold different value for persons with different goals. For example, those with aesthetic goals may have greater interest in whole muscle changes irrespective of whether changes occur as a result of contractile or non-contractile components increasing, whereas those with more performance-specific goals may have greater interest in fibre-specific adaptations or changes in muscle density. As such we believe that the different outcome methods, though both providing important information, ultimately provide different information and should be considered as such in interpretation.

4. Muscle Function Adaptations

Muscle function is often measured as either strength, relative endurance (repetitions performed at a submaximal %1RM load) or absolute endurance (repetitions performed with an absolute submaximal load). The nature of testing mode for these can vary considerably including free weights, resistance machines, and isokinetic or isometric dynamometers. Publications from the American College of Sports Medicine (ACSM) have suggested that HL promote greater strength adaptations whereas LL may promote greater endurance adaptations (though it is not specified whether they refer to relative or absolute endurance) [3,4].

However, these claims have received criticism [32,33] and authors of more recent reviews have reported similar increases in strength and absolute endurance adaptation irrespective of training load [2,10,34]. The similar changes in strength and absolute endurance have been suggested as possibly due to the inherent relationship between the two outcomes [35,36]. With this in mind it is important to consider the nature of the measures of muscular function employed in studies considering HL and LL training.

The recent meta-analysis referred to above [6] also examined a muscle function outcome (strength), again reporting no significant difference between HL and LL but a greater effect size in the HL condition ((mean ±SD) LL= 1.23 ±0.43; HL= 2.30 ±0.43). However, again some studies have utilised differing methods of measuring muscle function within their designs. For example, Mitchell et al. [19] reported a number of different muscle function related outcomes including strength (1RM and isometric maximal voluntary contractions) and relative endurance (repetitions to failure with both 30%- and 80%1RM loads in addition to total work). These varied with regards to whether changes significantly favoured the HL group (1RM and total work with 80%1RM) or the LL group (number of repetitions with 30%1RM). The authors of a more recent publication reported significantly greater strength adaptations for the back squat, but not bench press, 1RM when using 70-80% 1RM compared to 30-50% 1RM (although larger ESs for bench press were noted for the higher load group) [37]. Further, changes in relative endurance (repetitions to failure using 50%1RM) were significantly greater for the LL group. Interestingly, there were no significant between-group (HL vs. LL) differences for hypertrophy of the elbow flexors, extensors and quadriceps muscles. In contrast, the same group of authors reported significantly greater increases in 1RM for bench press, but not back squat, when training with 3RM compared to 10RM [38]. Another paper included in the above noted meta-analysis (Ogasawara et al., 2013) found no difference in elbow extension isokinetic strength between HL and LL groups

but did for bench 1RM. As with studies included in the hypertrophy component of this meta-analysis it is not clear how different outcomes were handled for these studies [19; Ogasawara et al. 2013] and, for reasons described below, this may have similarly impacted the ESs in favour of HL conditions.

It is interesting to consider the reasons for the divergent results within these studies and to consider the testing modes employed. We propose that one reason as to why there might be differing strength and hypertrophic adaptations might be that of skill specificity in motor recruitment (Behm & Sale, 1993). Motor control research suggests that a motor schema is highly specific to the task being practised [39,40], and though it could be argued that the higher number of repetitions associated with LL training could suggest a greater volume of practice favouring those conditions, motor schemata have also been reported to be *load/force* specific [41]. With this in mind, lifting a heavier load in a particular movement might serve to practise and refine that schema as a skill which would include the maximal synchronous recruitment of motor units and muscle fibres. This is a key reason why most maximal testing protocols include some sort of familiarisation or practice component within exercise science research [42]. Indeed the results of Mitchell et al. [19] support this contention: though the HL group had a greater increase in 1RM, possibly due to the motor schema refinement that likely occurred from training closer to their maximal load, there were no differences between the HL and LL groups for peak isometric maximal voluntary contraction, maximal power output, or rate of force development. The tendency for greater strength gains in the HL groups in the studies by Schoenfeld et al. [37,38] may also be due to this specificity of motor schema refinement. Further, the 1RM tasks measured were compound free weight movements (squat and bench press) which have been shown to require multiple (~3-5) familiarisation sessions even in moderately trained persons due to continued increases in 1RM [43] and improvements during these are likely attributable to neural and

learning effects [44]. In support of this are the results from Ogasawara et al. (2013) who reported significantly greater gains in bench press 1RM for the HL group however found no differences between groups for elbow extension strength. Thus, in the studies mentioned the apparent superiority of HL in enhancing strength may simply reflect better learning of the specific skills involved in the testing. In contrast, more simple strength tasks such as dynamometry of isolated joint movements require less refinement of motor schemata evidenced by the requirement for only a single familiarisation session to achieve reliable results [45,46]. However, that single familiarisation session is still essential to achieve valid results and therefore even with such simple tasks there is clearly a skill learning element to testing results. In our opinion researchers should therefore bear the specificity principle in mind when comparing the results of different training protocols, as the similarity of training and testing protocols is likely a key factor.

5. Exertion and discomfort

We also speculate that a secondary reason as to the differing results in these studies [19,37,38], particularly with respect to the changes in relative endurance, may relate to exertion and associated discomfort. The differentiation between perceptions of effort and discomfort have been highlighted recently as important [47], particularly within RT [48], for good reason.

Shimano et al. [49] considered rating of perceived exertion (RPE) values in trained and untrained persons performing a single set to momentary failure at 60%, 80% and 90% 1RM for back squat, bench press and arm curl. The authors reported no significant differences in RPE between load and exercise performed, with the exception of a significantly higher exertion for the back squat at 60% 1RM in trained persons ((mean ±SD)

8.8 ±0.7 vs. 6.9 ±1.9). This might suggest that the volume of repetitions preceding momentary failure may have produced a greater degree of discomfort resulting in a higher RPE value. Indeed further research has shown that when performing multiple sets to momentary failure, mean (±SD) RPE increases significantly from set one (50%1RM = 7.40±1.96 vs. 70%1RM = 7.73±1.44), to two (50%1RM = 8.60±0.99 vs. 70%1RM = 8.73±0.80), to three (50%1RM = 9.33±0.82 vs. 70%1RM = 9.47±0.74) with no difference between different loads [50]. We have quite specifically termed this discomfort rather than exertion for the following reason. The authors of these studies reported that participants exercised to momentary failure with verbal encouragement to ensure adequate motivation and effort, and RPE was measured using a Borg CR10 scale [51], where a value of 10 indicates maximal effort. In this case, each trial, irrespective of exercise, load, or training status should have resulted in a maximal value for effort since persons were exercising to momentary failure. Since participants did not report maximal values we can only assume that the participants were unclear as to how to report their perception of effort, and as such, potentially expressed their feelings of discomfort. Again, despite also using the Borg CR10 RPE scale and having participants train to momentary failure, Pritchett et al. [52] also reported RPE values of less than 10 for both acute and session RPE. However, RPE was significantly higher for the 60%1RM condition compared with 90%1RM, suggesting the LL with a higher number of repetitions incurred a higher discomfort than training at a HL. Based on this we hypothesise that people might find it more difficult to reach momentary failure with a LL because of a higher discomfort. As such, studies comparing HL and LL training where participants are said to have trained to momentary failure might be limited by a high discomfort in the LL group, preventing participants from reaching true momentary failure. We propose that in comparison of HL and LL groups the conduct of reaching momentary failure becomes all the more important in a LL group to maximally, sequentially recruit all

possible motor units. However, we should acknowledge that at present there are insufficient studies comparing LL training to momentary failure and not to momentary failure to determine how much of a meaningful difference a final repetition (e.g. reaching 'true' momentary failure) might make towards chronic adaptations.

6. Conclusion

When considering the findings of studies comparing the effects of HL and LL, there are a number of important factors to consider. These include the different outcomes related to morphological changes providing differing information, skill associated with the testing mode chosen (both load and task), and other psychosocial factors such as discomfort. We contend that different testing modes evidently reflect different outcomes and indeed they may hold different values for persons with different goals. Again it is possible that HL or LL may favour certain outcomes and not impact upon others. For example, if solely wishing to improve maximal strength of a specific task (such as a powerlifter wishing to improve back squat, deadlift or bench press) a recommendation might be to perform these specific exercises using heavy loads to attempt to catalyse both morphological and neural adaptations [30]. Whereas those more interested in improving muscular force production for health parameters or in a way that might be widely transferable may be able to utilise a variety of loading schemes [17].

We hope that the present piece has catalysed a more open mind-set toward some of the factors that must be considered with regards to interpretation of studies examining HL and LL in resistance training. The discussion of resistance training load is pertinent since most strength coaches first consider maximal strength testing to then make training recommendations based on % 1RM. The purpose of this piece is not necessarily to challenge others' recommendations regarding this topic; rather, we hope to provide practitioners with

the necessary understanding to interpret presently existing research on the topic and recommendations surrounding it that may on the surface seem to be contradictory. The impact of load in resistance training may produce differential adaptations in different aspects of morphology or function. Thus persons should first consider their desired training goals and then decide whether evidence would appear to suggest that the manipulation of load might impact those goals differentially. If the effect of load is presently equivocal for a particular outcome, there are potentially numerous practical implications of being able to self-select an external load. These include: reducing the need for specific facility memberships (e.g. where specifically heavy loads are available), motivating older persons or those who might be less confident using heavy loads, and allowing people to undertake home- or field-based resistance training intervention strategies. Ultimately these might serve to improve exercise adherence. As a final caveat to the content discussed, we recognise that there is very likely a threshold (below which would not produce continued recruitment because of recovery capacity of utilised motor units and muscle fibres, and thus would prevent someone ever reaching true momentary failure) which if not exceeded might produce sub-optimal adaptations, however this has not been identified empirically in any literature and is likely very individual to a person, and possibly exercise based on individual mechanics and muscle fibre type.

Compliance with ethical standards

Funding

No sources of funding were used to assist in the preparation of this article.

Conflicts of Interest

James Fisher, James Steele and Dave Smith declare that they have no conflicts of interest relevant to the content of this article.

References

1. Fisher J, Steele J, Smith D. Evidence-based resistance training recommendations for muscular hypertrophy. Med Sport. 2013;17(4):217-235.
2. Fisher J, Steele J, Bruce-Low S, et al. Evidence-based resistance training recommendations. Med Sport. 2011; 15(3):147-162.
3. Kraemer WJ, Adams K, Cafarelli E, et al. Progression models in resistance training for healthy adults. Med Sci Sports Exerc. 2002;34:364-80.
4. Ratamess NA, Alvar BA, Evetoch TK, et al. Progression models in resistance training for healthy adults. Med Sci Sports Exerc. 2009;41(3):687-708.
5. Schoenfeld B. Is there a minimum intensity threshold for resistance training-induced hypertrophic adaptations? Sports Med. 2013;43(12):1279-1288.
6. Schoenfeld BJ, Wilson JM, Lowery RP, et al. Muscular adaptations in low-versus high-load resistance training: A meta-analysis. Eur J Sport Sci. 2014;20:1-10, Epub ahead of print, doi:10.1080/17461391.2014.989922.
7. Schoenfeld BJ, Contreras B, Willardson JM, et al. Muscle activation during low- versus high-load resistance training in well-trained men. Eur J Appl Physiol. 2014;114(12):2491-2497.
8. Jenkins ND, Housh TJ, Bergstrom HC, et al. Muscle activation during three sets to failure at 80 vs. 30 % 1RM resistance exercise. Eur J Appl Physiol. 2015;115(11):2335-2347.
9. Looney DP, Kraemer WJ, Joseph MF, et al. Electromyographical and perceptual responses to different resistance intensities in a squat protocol: does performing sets to failure with light loads produce the same activity? J Strength Cond Res. 2016;30(3):729-799.

372 10. Denny-Brown D, Pennybacker JB. Fibrillation and fasciculation in voluntary muscle.
373 Brain. 1938;61(3):311-312.
374 11. Carpinelli R. The size principle and a critical analysis of the unsubstantiated heavier-
375 is-better recommendation for resistance training. J Exerc Sci Fit. 2008;6:67-86.
376 12. Enoka RM, Duchateau J. Inappropriate interpretation of surface EMG signals and
377 muscle fiber characteristics impedes progress on understanding the control of
378 neuromuscular function. J Appl Physiol. 2015;Epub ahead of print, doi:
379 10.1152/japplphysiol.00280.2015.
380 13. Adam A, De Luca CJ. Recruitment order of motor units in human vastus lateralis
381 muscle is maintained during fatiguing contractions. J Neurophysiol. 2003;90:2919-
382 2927.
383 14. Westad C, Westgaard RH, De Luca CJ. Motor unit recruitment and derecruitment
384 induced by brief increase in contraction amplitude of the human trapezius muscle. J
385 Physiol. 2003;552:645-656.
386 15. Petrofsky JS, Phillips CA. Discharge characteristics of motor units and the surface
387 EMG during fatiguing isometric contractions at submaximal tensions. Aviat Space
388 Environ Med. 1985;56:581-586.
389 16. Behm DG. Force maintenance with submaximal fatiguing contractions. Can J Appl
390 Physiol. 2004;29(3):274-290.
391 17. Garland SJ, Gossen R. The muscular wisdom hypothesis in human muscle fatigue.
392 Exerc Sport Sci Rev. 2002;30(1):45-49.
393 18. Boe SG, Stashuk DW, Doherty TJ. Motor unit number estimation by decomposition-
394 enhanced spike-triggered averaging: control data, test-retest reliability, and contractile
395 level effects. Muscle Nerve. 2004;29:693-699.

19. Hodson-Tole EF, Wakeling JM. Variations in motor unit recruitment patterns occur within and between muscles in the running rat (Rattus norvegicus). J Exp Biol. 2007;210:2333-2345.
20. Steele J, Fisher J. Scientific rigour: a heavy or light load to carry? Sports Med. 2014;44(1):141-142.
21. Schoenfeld B. Author's reply to Steele and Fisher: ''Scientific rigour: a heavy or light load to carry?": the importance of maintaining objectivity in drawing evidence-based conclusions. Sports Med. 2014;44:143-145.
22. Mitchell C J, Churchward-Venne TA, West DW, et al. Resistance exercise load does not determine training-mediated hypertrophic gains in young men. J Appl Physiol. 2012;113(1):71-77.
23. McCall GE, Byrnes WC, Dickinson A, et al. Muscle fiber hypertrophy, hyperplasia, and capillary density in college men after resistance training. J Appl Physiol. 1996;81(5):2004-2012.
24. Ogborn D, Schoenfeld BJ. The role of fiber types in muscle hypertrophy: implications for loading strategies. Strength Cond J. 2014;36(2):20-25
25. Vinogradova OL, Popov DV, Netreba AI, et al. Optimization of training: New developments in safe strength training. Hum Physiol. 2013;39(5):511-523
26. Netreba A, Popov D, Bravyy Yam et al. Responses of knee extensor muscles to leg press training of various types in human. Russian J Physiol. 2013;99(3):406-416
27. Hather BM, Adams GR, Tesch PE, et al. Skeletal muscle responses to lower limb suspension in humans. J Appl Physiol. 1992;72(4):1493-1498
28. Oates BR, Glover EI, West DW, et al. Low-volume resistance exercise attenuates the decline in strength and muscle mass associated with immobilization. Muscle Nerve. 2010;42(4):539-546

421 29. Adams GR, Bamman MM. Characterization and regulation of mechanical loading-
422 induced compensatory muscle hypertrophy. Compr Physiol. 2012;2(4):2829-2871.
423 30. Verdijk LB, Gleeson BG, Jonkers RAM, et al. Skeletal muscle hypertrophy following
424 resistance training is accompanied by a fiber-type specific increase in satellite cell
425 content in elderly men. J Gerontol. 2009;64A(3):332-339
426 31. Burd NA, Moore DR, Mitchell CJ, et al. Big claims for big weights but with little
427 evidence. Eur J Appl Physiol. 2013;113(1):267-267 .
428 32. Schuenke MD, Herman J, Staron RS. Preponderance of evidence proves "big"
429 weights optimize hypertrophic and strength adaptations. Eur J Appl Physiol. 2013;
430 113(1):269-271.
431 33. Wakahara T, Fukutani A, Kawakami Y, et al. Nonuniform muscle hypertrophy: Its
432 relation to muscle activation in training session. Med Sci Sports Exerc.
433 2013;45(11):2158-65.
434 34. Lexell J, Taylor CC. A morphometrical comparison of right and left whole human
435 vastus lateralis muscle: how to reduce sampling errors in biopsy techniques. Clin
436 Physiol. 1991;11(3):271–276.
437 35. Lemon PWR, Tarnopolsky MA, MacDougall JD, et al. Protein requirements and
438 muscle mass/strength changes during intensive training in novice bodybuilders. J
439 Appl Physiol. 1992;73(2):767-775.
440 36. Jones DA, Rutherford OM. Human muscle strength training: the effects of three
441 different regimes and the nature of resultant changes. J Physiol. 1987;391:1-11.
442 37. Carpinelli R, Otto RM, Winett RA. A critical analysis of the ACSM position stand on
443 resistance training: insufficient evidence to support recommended training protocols. J
444 Exerc Physiol. 2004;7:1-60.

445 38. Carpinelli R. Challenging the American College of Sports Medicine 2009 position
446 stand on resistance training. Med Sport. 2009;13:131-137.
447 39. Jungblut S. The correct interpretation of the size principle and its practical application
448 to resistance training. Med Sport. 2009;13(4):203-209.
449 40. Hickson RC, Hidaka K, Foster C. Skeletal muscle fibre-type, resistance training, and
450 strength-related performance. Med Sci Sports Exerc. 1994;26:593-598.
451 41. Mazzetti SA, Kraemer WJ, Volek JS, et al. The influence of direct supervision of
452 resistance training on strength performance. Med Sci Sports Exerc. 2000;32(6):1175-
453 1184.
454 42. Behm DG, Sale DG. Velocity specificity of resistance training. Sports Med.
455 1993;15(6):347-388.
456 43. Schoenfeld BJ, Peterson MD, Ogborn D, et al. Effects of low- versus high-load
457 resistance training on muscle strength and hypertrophy in well-trained men. J Strength
458 Cond Res. 2015;29(10):2954-2963.
459 44. Schoenfeld BJ, Ratamess NA, Peterson MD, et al. Effects of different volume-
460 equated resistance training loading strategies on muscular adaptations in well-trained
461 men. J Strength Cond Res. 2014;28(10):2909-2918.
462 45. Drowatzky JN, Zuccato FC. Interrelationships between selected measures of static
463 and dynamic balance. Res Q. 1967;38:509-10.
464 46. Mount J. Effect of practice of a throwing skill in one body position on performance of
465 the skill in an alternate position. Percept Mot Skills. 1996;83:723-32.
466 47. Schmidt, RA. Motor schema theory after 27 years: reflections and implications for a
467 new theory. Res Q Exerc Sport. 2003;74(4):366-375.
468 48. Brown LE, Weir JP. ASEP procedures recommendation I: accurate assessment of
469 muscular strength and power. J Exerc Physiol. 2001;4(3):1-21.

49. Soares-Caldeira LF, Ritt-Dias RM, Okuno NM, et al. Familiarization indexes in sessions of 1-RM tests in adult women. J Strength Cond Res. 2009;23(7):2039-2045.
50. Cronin JB, Henderson ME. Maximal strength and power assessment in novice weight trainers. J Strength Cond Res. 2007;18(1):48-52.
51. Graves JE, Pollock ML, Carpenter DM, et al. Quantitative assessment of full range-of-motion isometric lumbar extension strength. Spine. 1990;15(4):289-294.
52. Welsch MA, Williams PA, Pollock ML, et al. Quantification of full-range-of-motion unilateral and bilateral knee flexion and extension torque ratios. Arch Phys Med Rehabil. 1998;79(8):971-978.
53. Marcora S. Perception of effort during exercise is independent of afferent feedback from the skeletal muscles, heart, and lungs. J Appl Physiol. 2009;106:2060-2062.
54. Steele J. Intensity; in-ten-si-ty; noun. 1. Often used ambiguously within resistance training. 2. Is it time to drop the term altogether? Br J Sports Med. 2014;48(22):1586-1588.
55. Shimano T, Kraemer WJ, Spiering BA, et al. Relationship between the number of repetitions and selected percentages of one repetition maximum in free weight exercises in trained and untrained men. J Strength Cond Res. 2006;20:819-23.
56. Silva VL, Azevedo AP, Cordeiro JP, et al. Effects of exercise intensity on perceived exertion during multiple sets of bench press to volitional failure. J Trainol. 2014;3:41-46.
57. Borg GA. Psychophysical bases of perceived exertion. Med Sci Sports Exerc. 1982;14:377-81.
58. Pritchett RC, Green JM, Wickwire PJ, et al. Acute and session RPE responses during resistance training: Bouts to failure at 60% and 90% of 1RM. S Afr J Sports Med. 2009;21(1):23-26.

Figure 1 Adapted image of Forest plot by Schoenfeld, et al. [6]



Dotted line represents overall ES. Studies to the right of the dotted line used *in vitro* methods of measuring hypertrophy where studies to the left used *in vivo* methods of measuring hypertrophy.